

# Immunization Requirements for Childcare/Preschool

By Age 3 mo.	By Age 5 mo.	By Age 7 mo.	By Age 12 mo.	By Age 16 mo.	By Age 19 mo.	By Age 4 yrs.	By Age 5 yrs.
<a href="#">DTaP</a> 1 dose	<a href="#">DTaP</a> 2 doses	<a href="#">DTaP</a> 3 doses	<a href="#">DTaP</a> 3 doses	<a href="#">DTaP</a> 4 doses	<a href="#">DTaP</a> 4 doses	* <a href="#">DTaP</a> 4 doses	* <a href="#">DTaP</a> 5 doses
<a href="#">Polio (IPV)</a> 1 dose	<a href="#">Polio (IPV)</a> 2 doses	<a href="#">Polio (IPV)</a> 2 doses	<a href="#">Polio (IPV)</a> 2 doses	<a href="#">Polio (IPV)</a> 2 doses	<a href="#">Polio (IPV)</a> 3 doses	* <a href="#">Polio (IPV)</a> 3 or 4 doses	* <a href="#">Polio (IPV)</a> 3 or 4 doses
<a href="#">Hib</a> 1 dose	<a href="#">Hib</a> 2 dose	<a href="#">Hib</a> 2 doses	* <a href="#">Hib</a> 3 doses	* <a href="#">Hib</a> 3 or 4 doses	* <a href="#">Hib</a> 3 or 4 doses	* <a href="#">Hib</a> 3 or 4 doses	<a href="#">Hib</a> – no requirement
<a href="#">Hep B</a> 1 dose	<a href="#">Hep B</a> 2 doses	<a href="#">Hep B</a> 2 doses	<a href="#">Hep B</a> 2 doses	<a href="#">Hep B</a> 2 doses	<a href="#">Hep B</a> 3 doses	<a href="#">Hep B</a> 3 doses	<a href="#">Hep B</a> 3 doses
<a href="#">Pneumococcal (PCV)</a> 1 dose	<a href="#">Pneumococcal (PCV)</a> 2 doses	* <a href="#">Pneumococcal (PCV)</a> 3 doses	* <a href="#">Pneumococcal (PCV)</a> 4 doses	* <a href="#">Pneumococcal (PCV)</a> 4 doses	* <a href="#">Pneumococcal (PCV)</a> 4 doses	* <a href="#">Pneumococcal (PCV)</a> 4 doses	* <a href="#">Pneumococcal (PCV)</a> – no requirement
				<a href="#">MMR</a> 1 dose	<a href="#">MMR</a> 1 dose	<a href="#">MMR</a> 2 doses	<a href="#">MMR</a> 2 doses
				<a href="#">Varicella</a> 1 dose	<a href="#">Varicella</a> 1 dose	<a href="#">Varicella</a> 2 doses	<a href="#">Varicella</a> 2 doses

\*DTaP-Fifth dose not needed if dose #4 given after 4<sup>th</sup> B-day and 6 months from last dose

\*Polio-Fourth dose not needed if dose #3 given after 4<sup>th</sup> B-day and 6 months from last dose

\*Hib- May have fewer doses if start series late; last dose must be on or after 12 months of age; not required after age 5 years

\*PCV- May have fewer doses if start series late; last dose must be on or after 12 months of age; not required after age 5 years

\*Varicella-no doses needed if history of **diagnosed** Chickenpox disease